

Exercise 10.2

In Q. 1 to 3, choose the correct option and give justification.

1. From a point Q, the length of the tangent to a circle is 24 cm and the distance of Q from the centre is 25 cm. The radius of the circle is


(A) 7 cm (B) 12 cm (C) 15 cm (D) 24.5 cm.

Sol.  $r = \sqrt{(25)^2 - (24)^2}$  cm = 7 cm.

Option (A) is correct.

2. In figure, if TP and TQ are the two tangents to a circle with centre O so that  $\angle POQ = 110^\circ$ , then  $\angle PTQ$  is equal to

(A)  $60^\circ$  (B)  $70^\circ$   
(C)  $80^\circ$  (D)  $90^\circ$ .



Sol.  $\because$  TQ and TP are tangents to a circle with centre O.

such that  $\angle POQ = 110^\circ$

$\therefore OP \perp PT$  and  $OQ \perp QT$

$\Rightarrow \angle OPT = 90^\circ$  and  $\angle OQT = 90^\circ$

Now, in the quadrilateral TPOQ, we get

$$\therefore \angle PTQ + 90^\circ + 110^\circ + 90^\circ = 360^\circ$$

$$\Rightarrow \angle PTQ + 290^\circ = 360^\circ$$

$$\Rightarrow \angle PTQ = 360^\circ - 290^\circ = 70^\circ$$

Thus, the correct option is (B).

3. If tangents PA and PB from a point P to a circle with centre O are inclined to each other at angle of  $80^\circ$ , then  $\angle POA$  is equal to

(A)  $50^\circ$  (B)  $60^\circ$  (C)  $70^\circ$  (D)  $80^\circ$ .

Sol.  $\angle POA = \frac{1}{2} \angle AOB = \frac{1}{2} \times 100^\circ = 50^\circ$ .

Option (A) is correct.

4. Prove that the tangents drawn at the ends of a diameter of a circle are parallel.


Sol. In the figure, we have:

PQ is diameter of the given circle and O is its centre.

Let tangents AB and CD be drawn at the end points of the diameter PQ.

Since the tangent at a point to a circle is perpendicular to the radius through the point.

$$\therefore PQ \perp AB \Rightarrow \angle APQ = 90^\circ$$



And  $PQ \perp CD \Rightarrow \angle P Q D = 90^\circ$   
 $\Rightarrow \angle A P Q = \angle P Q D$

But they form a pair of alternate angles.

$\therefore AB \parallel CD.$

5. Prove that the perpendicular at the point of contact to the tangent to a circle passes through the centre.


Sol. Let perpendicular at the point of contact to the tangent does not pass through at centre.

$O'P \perp PT$  ... (i) [Given]

Join OP. As OP is radius.

$\therefore OP \perp PT$  ... (ii)

[Radius is perpendicular to tangent at the point of contact]



From (i) and (ii), we get

OP and O'P are perpendicular to PT

$\Rightarrow$  OP and O'P must coincide.


As only one perpendicular can be drawn from a point on a line.

Hence perpendicular from the point of contact, passes through the centre.

6. The length of a tangent from a point A at distance 5 cm from the centre of the circle is 4 cm. Find the radius of the circle.

Sol.  $OT = \sqrt{5^2 - 4^2}$

cm =  $\sqrt{9}$  cm = 3 cm.



7. Two concentric circles are of radii 5 cm and 3 cm. Find the length of the chord of the larger circle which touches the smaller circle.

Sol. OA = 5 cm, OT = 3 cm

Also  $OT \perp AB,$

Therefore,  $AT = \sqrt{25 - 9}$  cm = 4 cm


$\therefore AB = 2AT = 8$  cm.



## NCERT SOLUTIONS

8. A quadrilateral  $ABCD$  is drawn to circumscribe a circle (see figure). Prove that

$$AB + CD = AD + BC$$




**Sol.** We have  $AS = AP$ ;  $BP = BQ$ ;  $CQ = CR$  and  $DR = DS$ .

$$\begin{aligned} \text{Consider } AB + CD &= (AP + PB) + (CR + RD) \\ &= AS + BQ + CQ + DS \\ &= (AS + DS) + (BQ + CQ) \\ &= AD + BC. \end{aligned}$$

**Hence proved.**


9. In figure,  $XY$  and  $X'Y'$  are two parallel tangents to a circle with centre  $O$  and another tangent  $AB$  with point of contact  $C$  intersecting  $XY$  at  $A$  and  $X'Y'$  at  $B$ . Prove that  $\angle AOB = 90^\circ$ .



**Sol. Given:** A circle with centre  $O$  has three tangents  $XY$ ,  $X'Y'$  and  $AB$  at the points  $P$ ,  $Q$  and  $C$  respectively. Also  $XY \parallel X'Y'$ .

**To prove:**  $\angle AOB = 90^\circ$

**Construction:** Join  $OC$ ,  $OP$ ,  $OQ$ .



**Proof:** In  $\triangle AOP$  and  $\triangle AOC$ ,

$$PA = PC \quad \text{[Two tangents drawn from a point outside the circle]}$$

$$PO = CO \quad \text{[Radii of same circle]}$$

$$AO = AO \quad \text{[Common]}$$

Therefore,  $\triangle AOP \cong \triangle AOC$  [SSS criterion]

$$\therefore \angle 1 = \angle 2 \quad \dots(i)$$

Similarly, we can prove that

$$\angle 3 = \angle 4 \quad \dots(ii)$$

Now,  $\angle PAB + \angle QBA = 180^\circ$  [Sum of interior angles on the same side of transversal]

$$\Rightarrow 2\angle 2 + 2\angle 3 = 180^\circ \quad \text{[Using (i) and (ii)]}$$

$$\Rightarrow \angle 2 + \angle 3 = 90^\circ \quad \text{[Sum of angles of a triangle is } 180^\circ]$$

$$\Rightarrow \angle AOB = 90^\circ. \quad \text{Hence proved.}$$

10. Prove that the angle between the two tangents drawn from an external point to a circle is supplementary to the angle subtended by the line-segment joining the points of contact at the centre.


Sol.  $\angle OAP = \angle OBP = 90^\circ \dots(i)$

Also  $\angle AOB + \angle OBP + \angle BPA + \angle OAP = 360^\circ$

[Sum of angles of a quadrilateral is  $360^\circ$ ]

$$\Rightarrow \angle AOB + 90^\circ + \angle BPA + 90^\circ = 360^\circ$$

$$\Rightarrow \angle AOB + \angle BPA = 180^\circ.$$



11. Prove that the parallelogram circumscribing a circle is a rhombus.

Sol.  $AB = CD$  and  $BC = DA \dots(i)$


$$\begin{aligned} AB + CD &= AP + BP + CR + DR \\ &= AS + BQ + CQ + DS \\ &= AD + BC \end{aligned}$$

$$\Rightarrow 2AB = 2AD$$


$$\Rightarrow AB = AD$$

As adjacent sides of a parallelogram are equal.

$\therefore$  Parallelogram is a rhombus.




12. A triangle ABC is drawn to circumscribe a circle of radius 4 cm such that the segments BD and DC into which BC is divided by the point of contact D are of



lengths 8 cm and 6 cm respectively (see figure). Find the sides AB and AC.

Sol. Let the circumcircle touches AB and AC at E and F respectively.

Join OA, OB, OC, OE and OF.



We have

$$OD = 4 \text{ cm} = OE = OF \quad \text{[Radii]}$$

$$\text{Also } CD = 6 \text{ cm} = CF, \quad BD = 8 \text{ cm} = BE$$

$$\text{and } AE = AF = x \text{ cm} \quad \text{[Say]}$$

$\therefore$  The lengths of the two tangents from an external point to a circle are equal]

From figure,

$$ar(\Delta ABC) = ar(\Delta OAB) + ar(\Delta OBC) + ar(\Delta OCA) \dots(i)$$

But  $ar(\Delta ABC) = \sqrt{s(s-a)(s-b)(s-c)}$

$$= \sqrt{(14+x)(14+x-8-x)(14+x-14)(14+x-6-x)}$$

$$\left[ s = \frac{AB+BC+CA}{2} = \frac{8+x+8+6+6+x}{2} = 14+x \right]$$

$$= \sqrt{(14+x)6 \times x \times 8} = \sqrt{48x(14+x)}$$

$$ar(\Delta OAB) = \frac{1}{2} \times (8+x) \times 4 = 16+2x$$

$$ar(\Delta OBC) = \frac{1}{2} (6+8) \times 4 = 28$$

and  $ar(\Delta OCA) = \frac{1}{2} (6+x) \times 4 = 12+2x$

Now, putting these values in eqn. (i), we have

$$\sqrt{48x(14+x)} = (16+2x) + (28) + (12+2x)$$

$$\Rightarrow \sqrt{48x(14+x)} = 56+4x = 4(14+x)$$

$$\Rightarrow 48x(14+x) = 16(14+x)^2 \quad [\text{Squaring both sides}]$$

$$\Rightarrow 48x(14+x) - 16(14+x)^2 = 0$$

$$\Rightarrow 16(14+x)(3x-14-x) = 0$$

$$\Rightarrow 16(14+x)(2x-14) = 0$$

$$\Rightarrow x = -14, 7$$

Ignoring  $x = -14$  because length cannot be negative.

$$\therefore x = 7 \text{ cm}$$

Hence,  $AB = BE + AE = 8 + x = 8 + 7 = 15 \text{ cm}$

and  $AC = CF + AF = 6 + x = 6 + 7 = 13 \text{ cm}$ .

13. Prove that opposite sides of a quadrilateral circumscribing a circle subtend supplementary angles at the centre of the circle.

Sol.  $\Delta AOP \cong \Delta AOS$

[As  $AP = AS$ ,  $OP = OS$ ,  
 $AO$  is common]

$$\angle 1 = \angle 8$$

Similarly  $\angle 2 = \angle 3$

$$\angle 4 = \angle 5$$

$$\angle 6 = \angle 7$$

$$\Rightarrow \angle 1 + \angle 2 + \angle 3 + \angle 4 + \angle 5 + \angle 6 + \angle 7 + \angle 8 = 360^\circ$$

$$\Rightarrow 2\angle 1 + 2\angle 2 + 2\angle 6 + 2\angle 5 = 360^\circ$$

$$\Rightarrow (\angle 1 + \angle 2) + (\angle 6 + \angle 5) = 180^\circ$$

$$\Rightarrow \angle AOB + \angle COD = 180^\circ$$

Similarly, we can show

$$\angle AOD + \angle BOC = 180^\circ$$

